

Modal Fields of a Triangular Microcavity Laser with a Piercing Hole

Spiridonov A., Karchevskii E.

Kazan Federal University, 420008, Kremlevskaya 18, Kazan, Russia

Abstract

© 2018 IEEE. Eigenmodes of triangular microcavities with piercing holes in the center are computed as solutions of the Lasing Eigenvalue Problem using the system of Muller boundary integral equations and the Nyström method. The numerical study demonstrates that small centered holes in equilateral triangular microcavities can lead to a significant growth of the directionality of laser modes with the preservation of their low thresholds.

<http://dx.doi.org/10.1109/MMET.2018.8460276>

Keywords

lasing eigenvalue problem, microcavity laser, Muller boundary integral equation, Nyström method

References

- [1] M. Lebental, E. Bogomolny, and J. Zyss, "Organic micro-lasers: A new avenue onto wave chaos physics, " in A.B. Matsko (Ed.) Practical Applications of Microresonators in Optics and Photonics, CRC Press, Roca Baton, pp. 317-353, 2009.
- [2] E. I. Smotrova, A.I. Nosich, T.M. Benson, and P. Sewell, "Cold-cavity thresholds of microdisks with uniform and non-uniform gain: quasi-3D modeling with accurate 2D analysis, " IEEE J. Sel. Topics Quant. Electron., vol. 11, pp. 1135-1142, 2005.
- [3] E. I. Smotrova, V. Tsvirkun, I. Gozhyk, C. Lafargue, C. Ulysse, , et al., "Spectra, thresholds, and modal fields of a kite-shaped microcavity laser, " J. Opt. Soc. Am. B, vol. 30, no 6, pp. 1732-1742, 2013.
- [4] E. I. Smotrova and A. I. Nosich, "Mathematical study of the twodimensional lasing problem for the WG modes in a circular dielectric microcavity, " Opt. Quant. Electron, vol. 36. pp. 213-221, 2004.
- [5] O. V. Shapoval, K. Kobayashi, and A. I. Nosich, "Electromagnetic engineering of a single-mode nanolaser on a metal plasmonic strip placed into a circular quantum wire, " IEEE J. Sel. Topics Quant. Electron., vol. 23, no 6, art no 1501609, 2017.
- [6] E. Karchevskii and A. Nosich, "Methods of analytical regularization in the spectral theory of open waveguides, " Proc. Int. Conf. Math. Methods in EM Theory (MMET 2014), Dnipropetrovsk, pp. 39-45, 2014.
- [7] A. I. Nosich, "Method of analytical regularization in computational photonics, " Radio Science, vol. 51, no 8, pp. 1421-1430, 2016.
- [8] I. O. Sukharevsky, A. I. Nosich, and A. Altintas, "Manipulation of backscattering from a dielectric cylinder of triangular cross-section using the interplay of GO-like ray effects and resonances, " IEEE Trans. Antennas Propagat., vol. 63, no 5, pp. 2162-2169, 2015.
- [9] I. O. Sukharevsky, M. Lebental, B. Dietz, C. Lafargue, and S. Bittner, "Dielectric equilateral triangle microresonators: Integral equations and semiclassical physics approaches, " Proc. SPIE, vol. 10518, art. no 105181U-1, 2018.

- [10] A. O. Spiridonov, E. M. Karchevskii and A. I. Nosich, "Spectra, thresholds, and modal fields of a circular microcavity laser transforming into a square, " Proc. Int. Conf. Transp. Opt. Networks (ICTON-2015), art. no. 7193654, 2015.
- [11] A. O. Spiridonov and E. M. Karchevskii, "Field patterns of whisperinggallery and bow-Tie modes of elliptic microcavity laser with a circular active region, " Proc. Int. Conf. Transp. Opt. Networks (ICTON-2016), art. no. 7550639, 2016.
- [12] A.S. Zolotukhina, A. O. Spiridonov, E. M. Karchevskii, and A. I. Nosich "Lasing modes of a microdisk with a ring gain area and of an active microring, " Opt. Quant. Electron., vol. 47, no. 12, pp. 3883-3891, 2015.
- [13] A. S. Zolotukhina, A. O. Spiridonov, E. M. Karchevskii, and A. I. Nosich, "Electromagnetic analysis of optimal pumping of a microdisk laser with a ring electrode, " Appl. Phys. B. , vol. 123, art. no. 32, 2017.
- [14] A. O. Spiridonov, et al., "Symmetry accounting in the integral-equation analysis of the lasing eigenvalue problems for two-dimensional optical microcavities, " J. Opt. Soc. Am. B, vol. 34, pp. 1435-1443, 2017.
- [15] A. O. Spiridonov and E. M. Karchevskii, "Mathematical and numerical analysis of the spectral characteristics of dielectric microcavities with active regions, " Proc. Int. Conf. Days on Diffraction (DD-2016), pp. 390-395, 2016.
- [16] A. O. Spiridonov and E. M. Karchevskiy, "Projection methods for computation of spectral characteristics of weakly guiding optical waveguides, " Proc. Int. Conf. Days on Diffraction (DD-2013), pp. 131-135, 2013.
- [17] Y.-Z. Huang, Q. Chen, W.-H Guo, Q.-Y. Lu, and L.-J Yu, "Mode characteristics for equilateral triangle optical resonators, " IEEE J. Sel. Top. Quant. Electron., vol. 12, no 1, pp. 59-65, 2006.
- [18] C. Lafargue, M. Lebental, A. Grigis, C. Ulysse, I. Gozhyk, N. Djellali, J. Zyss, and S. Bittner, "Localized lasing modes of triangular organic microlasers, " Phys. Rev. E., vol. 90, pp. 052922/17, 2014.
- [19] J. Wiersig and M. Hentschel, "Unidirectional light emission from high-Q modes in optical microcavities, " Phys. Rev. A, vol. 73, no 3, pp. 031802(R)/4, 2006.
- [20] N. Djellali, I. Gozhyk, D. Owens, S. Lozenko, M. Lebental, et al. "Controlling the directional emission of holey organic microlasers, " Appl. Phys. Lett., vol. 95, no 10, pp. 101108/3, 2009.
- [21] W. G. Bickley, "Two-dimension potential problems concerning a single closed boundary, " Phil. Trans., vol. 228, pp 235-273, 1929.